

Knowledge, Attitude, and Practice towards Family Planning among Married Women in ALTI city, Alkamlin locality, AL-Gezira State, Sudan from February – March 2023

Abstract

Background: Researches have shown highest awareness but low utilization of contraceptives making the situation a serious challenge. Most of women in reproductive age group know little or have incorrect information about family planning methods. Even when they know the name of some of the contraceptives, they do not know where to get them or how to use it. These women have negative attitude about family planning, whereas some have heard false and misleading information, the current study aimed in assessing the knowledge, attitude, and practice of family planning among married women in ALTI city.

Justification: Considering the high risk of maternal morbidity and mortality, increased risks of unintended pregnancy, this study aims to explore the knowledge, attitude, and practice (KAP) of family planning (FP) and associated factors among married women living in ALTI city February – 2023.

Methods: Cross sectional study. 45 women were interviewed. Data were collected using a structured and pretested questionnaire, which included study participants' socio-demographic characteristics, access to FP services, knowledge, attitude, and practice of FP.

Study population: This study targeted married women in childbearing age in ALTI city February-2023.

Keywords

Family planning • Pregnancy • Reproductive age • Women • Fetus

Review Article

Farah Abdeljalil Elsiddig Dafalla*

Faculty of Medicine, University of Gezira, Sudan

***Correspondence:** Farah Abdeljalil Elsiddig Dafalla, Faculty of Medicine, University of Gezira, Sudan, Tel: 00249119293284; E-mail: farohgleel619@gmail.com

Received: 10 October, 2023; **Processed:** 06 November, 2023; **Accepted:** 13 November, 2023

Copyright: © 2023 Dafalla FAE. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Introduction

Family planning (FP) is defined as a way of thinking and living that is adopted voluntarily upon the bases of knowledge, attitude, and responsible decisions by individuals and couples [1]. Family planning refers to a conscious effort by a couple to limit or space the number of children they have with contraceptive methods [2].

Family planning deals with reproductive health of the mother, having adequate birth spacing, avoiding undesired pregnancies and abortions, preventing sexually transmitted diseases and improving the quality of life of mother, fetus and family as a whole [3,4].

The Federal Ministry of Health (FMOH) has undertaken many initiatives to reduce maternal mortality. Among these initiatives, the most important is the provision of family planning at all levels of the healthcare system [5,6].

Currently, short-term modern family planning methods are available at all levels of governmental and private health facilities, while long-term method is being provided in health centres, hospitals and private clinics [7-9].

Most of reproductive age women know little or incorrect information about family planning methods. Even when they know some names of contraceptives, they don't know where to get them or how to use it. These women have negative attitude about family planning, while some have heard false and misleading information [10,11]

Objectives

General objective

To assess knowledge, attitude, and practice toward family planning among married women in Alti town February – 2023.

Specific objectives

1. To assess the Concept of family planning among married women in Alti town.
2. To asses the attitude of married women toward family planning.
3. To assess the practice of family planning.
4. To know does the married women know all the methods of contraceptive.
5. To assess the knowledge of the complications from this contraception.

Literature Review

Previous study

Measuring family planning quality and its link with contraceptive use in public facilities in Burkina Faso, Ethiopia, Kenya and Uganda July 2018. Adolescents and long-acting reversible contraception: lessons from Mexico June 2018. Knowledge and acceptability of long-acting reversible contraception among adolescent women receiving school-based primary care services June 2018. Who's that girl? A qualitative analysis of adolescent girls' views on factors associated with teenage pregnancies

in Bolgatanga, Ghana June 2018. Effects of relationship context on contraceptive use among young women June 2018. Knowledge and use of emergency contraception among students of public secondary schools in Ilorin, Nigeria June 2018. Emergency contraception in Albania: A multi-methods qualitative study of awareness, knowledge, attitudes, and practices May 2018. Postpartum intrauterine device placement: a patient-friendly option May 2018. ACOG Committee Opinion No. 735: Adolescents and long-acting reversible contraception: implants and intrauterine devices May 2018. Married women's autonomy and post-delivery modern contraceptive use in the Democratic Republic of Congo April 2018. Effect of self-administration versus provider-administered injection of subcutaneous depot medroxyprogesterone acetate on continuation rates in Malawi: a randomised controlled trial April 2018. Factors associated with removal difficulties of etonogestrel-containing contraceptive implants (Nexplanon®) April 2018. Influence of age on tolerability, safety and effectiveness of subdermal contraceptive implants April 2018. Client satisfaction and decision making amongst females visiting family planning clinics in Hyderabad, Pakistan February 2018. Recommendations for contraception: examining the role of patients' age and race February 2018. Developing mHealth messages to promote postmenstrual regulation contraceptive use in Bangladesh: participatory interview study January 2018. Pill Magic – The Economist January 2018. Factors associated with non-adherence to instructions for using the Nestorone®/ethinyl estradiol contraceptive vaginal ring January 2018. Effects of immediate postpartum contraceptive counseling on long-acting reversible contraceptive use in adolescents December 2017. Knowledge and opinions of postpartum mothers about the lactational amenorrhea method: the Turkish experience December 2017 [12].

Methodology

Descriptive cross-sectional facility based study

45 women were interviewed. Data were collected using a structured and pretested questionnaire, which included study participants' socio-demographic characteristics, access to FP services, knowledge, attitude, and practice of FP.

Study population

This study targeted married women in childbearing age in ALTI town February-2023.

Study area: Alti rural hospital, Alti city, Gazira state, Sudan

Inclusion criteria

Married women in childbearing age in ALTI town.

Exclusion criteria

1. All women outside ALTI town.
2. All unmarried women.
3. Older women in menopause age.

Ethical approval

Verbal consent from selected samples without mentioning their name for keeping their privacy

Sample size

N= 450 10% of N =45 N=45

Sample technique

Quantities random technique

KAP study

I adopted it on old questionnaire and adjusted to my research

Knowledge

- 80% if true: Good knowledge
- 60% - 79%: Average knowledge
- Less than 60%: Poor knowledge

Attitude

- More than 50% positive
- Less than 50% negative

Practice

If answer all: good practice

If answer less than half: poor practice

Data collection



Structured questionnaire was used to interview the study population.





Data analyses and data presentation

The data were collected and analysed by Google form, software values were expressed in percentage and figures (Figures 1-20).

Limitation of this study

- Small numbers of study population
- Small coverage area with questioner









Results

The study showed that the overall proper knowledge, attitude and practice of women towards family planning (FP) was 63%, 40%, and 41% respectively. Factors associated with the practice of FP were residence, marital status, educational status, age, occupation, and knowledge, and attitude and number of children. In this study, the level of knowledge and attitude towards family planning was relatively low and the level of family planning

utilization was quite low in comparison with many studies. Every health worker should teach the community on family planning holistically to increase the awareness so that family planning utilization will be enhanced. Besides, more studies are needed in a thorough investigation of the different reasons affecting the non-utilizing of family planning and how these can be addressed are necessary.







Discussion

The results of the present study showed that 63% of study participants had good knowledge, 41% had favourable attitude, and 40% had good practice towards family planning.

It has also found that women who completed primary & secondary education were practicing family planning

than those who were uneducated (64% and 20.6%) respectively. This finding was in line with a study done in Jemma, Ethiopia [9]. This might be due to the fact that women who were able to read and write would think in which FP activities are useful to be economically, self-sufficient and more likely to acquire greater confidence and personal control in marital relationships including the discussion of family size and contraceptive use.



This study showed that, age of the study participants had an association with practicing FP. Those reproductive age women whose age >30 years were practicing family planning better than those whose age <18 years. This finding was in line with a study done in India. This might be due to the reason that, when age increases mothers awareness; attitude and practice towards family planning may increase. In addition, as age increases the chance of practicing sexual intercourse increases and as a result they would be interested to utilize family planning in one or another way.

The current study also showed that knowledge and attitude of reproductive age women were related to FP utilization. Those reproductive age women who had good knowledge were utilized FP better than from those who were less knowledgeable.

Those participants with favourable attitude were practicing better than those who had unfavourable attitude. This might be because knowledge and attitude for specific activities are the key factors to start behaving and maintaining it continuously.




Figure 15. This chart represents thinking of ALTI population about contraceptives that cause infertility. N= 45.




Figure 18. This chart represents the place where the woman take contraceptive. N= 45.




Figure 16. This chart represents thinking of take husband Permission first before using contraceptive methods. N= 45.




Figure 19. This chart represents if any side effects happen from contraceptive. N= 45.




Figure 17. This chart represents which type of contraceptive does the woman use. N= 45.




Figure 20. This chart represents the source of having this knowledge. N= 4.

Conclusion & Recommendation

In conclusion the study showed that married women in ALTI city where a marginalized group in terms of family planning and that their overall capacity for family planning, practices and professional practices were low. Contraception was unpopular among the women of Altay,

mainly due to the lack of educational qualifications and awareness of family planning. In addition, there were no family planning initiatives and campaigns in ALTI city. Therefore, strengthening family planning interventions and increasing access to basic health services and education is indispensable to improve maternal and child health.

References

1. Bernard, Roger Paul. "Factors governing IUD performance." *Am J Public Health* 61 (1971): 559-567.
2. MacCorquodale, Donald W. "Analysis of a family planning program in Guatemala." *Public Health Rep* 85 (1970): 570-574.
3. Maccorquodale, Donald W. "Characteristics of clinic personnel may influence family planning acceptance." *Health Serv Rep* 87 (1972): 782-786.
4. MacCorquodale, Donald W., and Thomas W. Pullum. "A mathematical model for determining effectiveness of family planning clinics." *Stud Fam Plann* 5 (1974): 232-238.
5. Reynolds, Jack. "Delivering family planning services: Autonomous vs. integrated clinics." *Fam Plann Perspect* 2 (1970): 15-22.
6. Kurup, R. S., P. S. G. Nair, and N. V. George. "IUCD acceptors in Trivandrum, Kerala, South India." *Soc Biol* 17 (1970): 226-230.
7. Sivin, Irving. "Fertility decline and contraceptive use in the international postpartum family planning program." *Stud Fam Plann* 12 (1971): 248-256.
8. McCalister, Donald, and Victor Thiessen. "Prediction in family planning. Prediction of the adoption and continued use of contraception." *Am J Public Health Nations Health* 60 (1970): 1372-1381.
9. MacCorquodale, Donald W. "Philippine physicians' influence on acceptors' selection of the IUD." *Stud Fam Plann* 5 (1974): 140-143.
10. Osteria, Trinidad S. "A cost-effectiveness analysis of family planning programs in the Philippines." *Stud Fam Plann* 4 (1973): 191-195.
11. MacCorquodale, Donald W. "Family planning education in the Philippines: Findings related to differences in training." *J Med Educ* 49 (1974): 686-690.
12. MacCorquodale, Donald W. "The attitudes of Philippine family planning physicians towards sterilization." *Soc Sci Med* 8 (1974): 591-593.

Citation: Dafalla, Farah Abdeljalil Elsiddig. "Knowledge, Attitude, and Practice towards Family Planning among Married Women in ALTI city, Alkamlin locality, AL-Gezira State, Sudan from February – March 2023." *J Gynecol Matern Health* 1 (2023): 105. DOI: [10.59462/JGMH.1.1.105](https://doi.org/10.59462/JGMH.1.1.105)